

Research Paper

Bioaccumulation of some metals in different organs of frozen Spotted seabass, Misurata-Libya

Esmail M. Alhemmal^{1*}, Adel O. Abudabos², Hanan M. Alfgih³, Hajer A. Alshitshat⁴, Jamal M. Almestiry⁵ and Halima A. Haded⁶

^{1,2,3,4,6}Department of Zoology, Faculty of Science, Misurata University, Misurata Libya.
⁵Department of Food Technology, Faculty of Agriculture, Misurata University, Misurata Libya.
Corresponding author E-mail: esmail74science@gmail.com

Received 21 February 2018; Accepted 28 March, 2018

This study aim to estimate bioaccumulation of some metals (Cr, Mn, Fe, Cu, Zn and Pb) in different fish tissue (skin, Muscle, liver and gills) of *Dicentrarchus punctatus*. The samples were collected from local supermarket in Misurata city, Libya in summer, 2017. Tissue of this study was analyzed by atomic absorption spectrophotometer (AAS) to find the concentration of heavy metal (mg/kg). The results showed that the mean concentration of Zn was higher than Pb in skin, gills. Furthermore, Fe, Zn and Mn were higher in muscle than liver, except Cu and Zn their concentrations

were higher in liver. While low concentration of Pb in all study organs. The order of metals in liver, gills, skin and muscle samples were found to be Cu > Fe > Mn > Zn > Cr = Pb; Fe > Zn > Cr > Mn > Cu > Pb; Fe > Zn > Mn > Cr = Cu > Pb respectively. Some levels of metals (Cr and Fe) in this study were higher than the maximum permissible limits of FAO and WHO for human consumption.

Keywords: AAS, bioaccumulation, organs, metals, spotted seabass

INTRODUCTION

Marine organisms could bioaccumulate numerous inorganic pollutants, with chromium (Cr), manganese (Mn), iron (Fe), copper (Cu), Zinc (Zn) and lead (Pb) contaminants being the most commonly studied elements. These elements are characterized by bioaccumulation, biomagnification in the food chain and tendency to persist in the environment (Morgano et al., 2014). Some aquatic organisms such as fish are often at within the aquatic food chain and large amount concentrate of metals in water that build up by drinking sea water during feeding and ion-exchange of dissolved metals across lipophilic membranes and absorption on tissue and membrane surface (Mendil et al., 2005; Agbozu et al., 2007). Heavy metals are taken up through different organs of fish because of the affinity between them, for this reason, many heavy metals are concentrated at different levels in different organs of the body (Rao and Padmaja, 2000; Bervoets et al., 2001; Karadede et al., 2004). Multiple factors including season,

physical and chemical properties of water can play a significant role in metal accumulation in different fish tissues (Hayat et al., 2007; Romeo et al., 1999). The intake rate of these heavy metals by humans through the consumption of fish causes serious health hazards (Luoma and Rainbow, 2008).

Some heavy metals such as Zn, Cu, Mn and Fe are essential for growth and well being of living organisms including man. However, they likely show toxic effects when organisms are exposed to levels higher than normally required. Other elements such as Pb are not essential for metabolic activities and exhibit toxic properties even with traces levels (FAO, 1992). Fish is a healthy and cheap protein source for a bigger percentage of world population (Hajeb et al., 2009). Fish is a qualified food in terms of energy and nutritious components (essential multi unsaturated fatty acids, exogenic amino acids, minerals and water-soluble vitamins) in human nutrition (Usydyus et al., 2009). This study aimed to

determine some essential and nonessential metals (Cr, Mn, Fe, Cu, Zn and Pb) in the skin, muscle, gills and liver of *Dicentrarchus punctatus*. Moreover, the permissible limit of these heavy metals via fish consumption was discussed.

MATERIALS AND METHODS

Sample preparation

Ten samples of spotted seabass (*Dicentrarchus punctatus*) were collected (frozen fish) from local supermarket in Misurata city, during summer, 2017 (Figure 1). Collected fish were transferred to the laboratory of Aquatic Biology unit, department of zoology, University of Misurata, Libya in poly-ethylene bags, put into container of polystyrene ice box. Samples were washed with distilled water, and dissected for skin, muscle (dorsal muscle without skin), gill and liver. 0.5 gram of the samples were put in volumetric flask, added with 10 mL of concentrated (56%, Merck) nitric acid overnight. The mixture heated on a hot plate in fume hood until a white fume was observed which shows that the digestion of tissue was complete (Tyokumbur, 2012). Sample was allowed to cool at room temperature, then filtered using filter paper into a 100 cm³ volumetric flask and made to mark with distilled water.

Sample analysis

Heavy metals analysis was measured by atomic absorption spectrophotometer (AAS) model number ITEM No. 19102,12, AAS, HiTACHI, of Central laboratory of analysis of Iron and steel factory, Misurata, Libya. Results are expressed as milligrams per gram.

Data statistics

Statistics were performed using SPSS 11.5 software. A P-value of 0.05 or less was considered statistically significant. For T-test ANOVA compared the metals concentration in skin, muscle, liver and gills.

RESULTS

The study was carried out to determine heavy metals concentration (Cr, Mn, Fe, Cu, Zn and Pb) in different organs such as skin, muscle, liver and gills of *D. punctatus*, collected from local supermarket in Misurata city. Table 1 shows the concentration mean of heavy metals in different organs of fish used for the study. The mean concentrations of Cr were found to be 0.16 ± 0.09 , 0.06 ± 0.00 , 0.02 ± 0.00 and 0.06 ± 0.00 mg/g in the skin,

muscle, liver and gills respectively. Manganese concentrations ranged from 0.03 ± 0.08 – 0.38 ± 0.15 mg/g in organs of fish (Table 1). The highest levels of Fe were found in gills (1.59 ± 0.21) with compared all organs of *D. punctatus*, while the highest levels of Cu were found in liver (0.96 ± 4.20) also Zn (1.45 ± 0.11) was found in skin. Table 1, shown that statistical analysis of metal concentrations showed a significant ($P < 0.05$) difference between gills and liver of Cr and Cu. The results of this study in (Figure 1) shows that skin had the highest value in Fe and Zn concentrations. Fe concentration is highest in muscle when compared with all organs of *D. punctatus*. The mean level of Copper (Cu) in liver was highest while the mean of Zinc (Zn) values were highest in gills. Mean level of Lead (Pb) in organs of fish were lower than others in the currently study.

DISCUSSION

Variation in levels of metals among the fish species is apparent possibly due to differences in metal concentrations in the local environment and chemical characteristics of water from which fishes were caught, ecological needs, metabolism and feeding patterns of fishes, the season in which studies were carried out and also length and weight of the fishes can play a role in accumulation of metals (Rauf et al., 2009). The present work revealed marked increase of heavy metals concentration such as Cr, Fe and Zn in skin, while muscle had high concentration of Fe. Also, Singh et al. (1991) and Yilmaz, (2003) suggested that, Skin tissue showed elevated levels of metals in comparison with muscle tissue. Muscle comparison skin does not come into direct contact with the metals as it is totally covered externally by the skin, that in many ways helps the fish to ward off the penetration of the trace metals and also it is not an active site for detoxification, and therefore transport of trace metals from other tissues to muscle (Dural et al., 2010). Furthermore, according to Bat et al., (2012) the skin of fishes may be an important site for the uptake of metals due to their high surface area to body ratio.

In the investigation, liver has showed higher concentrations of metals such as Cu and concentrations of Fe in gills can be influenced by absorption of metals onto the gill surface, as well as by formation of complexes between the metals and the mucous, which is often impossible to remove them from lamellae prior to the analysis (Erdoğrul and Erbilir, 2007). Furthermore, that iron metal (Fe) was used in anabolic of hemoglobin make the gill a prime site for Fe (essential metal) accumulation (Bat et al., 2012). The physiological role of the tissue in fish metabolism influences the concentration of metals. Some tissues such as the liver and the gills are metabolically active, as compared to low tissues metabolism like muscles (Marcovecchio et al., 1991) and

Table 1. Concentrations mean of heavy metals (mg/kg) in different organs in *D. punctatus*.

Sample	Concentration of heavy metals (mean±S.E.)					
	Cr	Mn	Fe	Cu	Zn	Pb
Skin	0.09±0.16	0.14±0.15	0.24±0.16	0.08±0.03	1.45±0.11	0.02±0.00
Muscle	0.06±0.00	0.16±0.32	0.44±0.24	0.02±0.00	0.39±0.04	0.02±0.00
Liver	0.02±0.00*	0.08±0.03	0.30±0.14	4.20±0.96*	0.21±0.50	0.02±0.00
Gills	0.06±0.00*	0.38±0.15	1.59±0.21	0.17±0.15*	0.86±0.29	0.02±0.00

Data are presented as means±S.E. of organs of *D. punctatus* of five replicates, (n=5). Values with different superscript letters (*) in the same column are significantly different at the 0.05 level (P≤0.05).




Figure 1. Distribution of metals in fish. A: Skin, B: Muscle, C: Liver, D, Gills.

protein metallothionein.

Finally, the observed differences can be explained by the fact that the concentrations of these metals depend to a great extent on the part of the fish analyzed (Tüzen, 2003). Moreover, ecological factors such as season, location, nutrient availability and temperature and salinity of the water, may be contribute to variations in the metal concentrations in fishes (Bashir et al., 2012).

ACKNOWLEDGEMENTS

Our sincere thanks to Mohamed Alshaafi for data analysis and the authors would also like to thank Mr. Mohamed Albaowr, staff of Central Laboratory of Analysis (CLA) of Iron and Steel Factory, Misurata, Libya,

for their help in heavy metal analysis using Atomic Absorption Spectrophotometer (AAS).

REFERENCES

Agbozu IE, Ekweozor IKE, Opuene K (2007). Survey of heavy metals in the catfish *Synodontis clarias*. *International Journal Environment of Science Technology*, 4 (1): 93-97.

Bashir MK, Steven S, Pandit R (2012). The determinants of rural household food security in the Punjab, Pakistan: an econometric analysis. Working paper 1203, School of Agricultural and Resource Economics, University of Western Australia, Crawley, Australia online available at <http://ageconsearch.umn.edu/bitstream/122526/2/working%20paper%201203.pdf>.

Bat L, Sezgin M, Üstün F, Şahin F (2012). Heavy metal concentrations in ten species of fishes caught in Sinop coastal waters of the Black Sea, Turkey. *Turkish Journal of Fisheries and Aquatic Sciences*,

- 12:371-376.
- Bervoets L, Blust R, Verheyen R (2001). Accumulation of metals in tissues of three spined stickleback (*Gasterosteus aculeatus*) from natural fresh waters. *Ecotoxicology and Environmental Safety*, 48 (2), 117 -27.
- Dural M, Goksu MZL, Ozak AA (2010). Investigation of heavy metal levels in economically important fish species captured from the Tuzla lagoon. *Food chemistry*, 27: 521–526.
- Erdoğrul O, Erbilir F (2007). Heavy metal and trace elements in various fish samples from Sir Dam Lake, Kahramanmaraş, Turkey, Environment monitoring. *Assess.* 130: 373–379.
- FAO (1992). Committee for inland fisheries of Africa. Report of the third assessing of the working party on pollution and fisheries Accar, Ghana, 25-29 November 1991-FAO. Fisheries report. No. 471.
- Hajeb P, Jinap S, Ismail A, Fatimah AB, Jamilah B, Rahim MA (2009). Assessment of mercury level in commonly consumed marine fishes in Malaysia, *Food Control*, 20: 79–84.
- Hayat S, Ali B, Ahmad A (2007). Salicylic acid: Biosynthesis, metabolism and physiological role in plants. In S. Hayat, A. Ahmad (eds.), *Salicylic acid: A plant hormone*, (pp. 1 – 14). Springer, Dordrecht, The Netherlands.
- Karadede H, Oymak SA, Unlu E (2004). Heavy metals in mullet, *Liza abu*, and cat fish, *Silurus triotegus*, from the Atatürk Dam Lake (Euphrates), Turkey. *Environmental International*, 30, 183 -188.
- Luoma SN, Rainbow PS (2008). *Metal contamination in aquatic environments: Science and lateral management*. Cambridge (UK): Cambridge Univ. p.573.
- Marcovecchio JE, Moreno VJ, Pérez A (1991). Metal accumulation in tissues of sharks from the Bahía Blanca Estuary, Argentina. *Marine Environmental Research*, 31(4): 263-274.
- Mendil M, Uluozlu OD, Hasdemir E, Tuzen M, Sari H, Suicmez M (2005). Determination of trace metal levels in seven fish species in lake in Tokat, Turkey, *Food Chemistry*, 90(1-2): 175-179.
- Morgano MA, Rabonato LC, Milani RF, Miyagusku L, Quintaes KD (2014). As, Cd, Cr, Pb and Hg in seafood species used for sashimi and evaluation of dietary exposure. *Food Control*, vol. 36, no. 1, pp. 24-39. <http://dx.doi.org/10.1016/j.foodcont.2013.07.036>.
- Rao LM, Padmaja G, (2000). Bioaccumulation of heavy metals in *M. Cyprinoids* from the harbor waters of Visakhapatnam. *Bull. Pure Applied Sci.*, 19: 77-85.
- Rauf A, Javed M, Ubaidullah M (2009). Heavy metal levels in three major carps (*Catla catla*, *Labeo rohita* and *Cirrhina mrigala*) from the River Ravi, Pakistan. *Pakistan Veterinary Journal*, 29(1) 24–26.
- Romeo M, Siau Y, Sidoumou Z, Gnassia-Barellia M, (1999). Heavy metal distribution in different fish species from the Mauritania coast. *Sci. Total Environ.*, 232: 169-175.
- Singh JG, Chang-Yen I, Stoute VA, Chatergoon L (1991). Distribution of selected heavy metals in skin and muscle of five tropical marine fishes. *Environmental Pollution*, 69(2-3): 203-215.
- Trachurus mediterraneus* from Iskenderum Bay, Turkey. *Environmental Research*, 92(3): 277-281.
- Tüzen M (2003). Determination of heavy metals in fish samples of the middle Black sea (Turkey) by graphite furnace atomic absorption spectrometry. *Food chemistry*. 80(1): 119-123.
- Tyokumbur ET (2012). Studies on physico-chemical parameters and cyclopoid copepod abundance in Awba Reservoir, Ibadan. *International Journal of Science and Development Studies*. Vol.7.No.7, 81-93.
- Usydus Z, Szlinder-Richert J, PolakJuszczak L, Komar K, Adamczyk M, Malesa-Cieciewicz M, Ruczynska W (2009). Fish products available in polish market assessment of the nutritive value and human exposure to dioxins and other contaminants, *Chemosphere*, 74: 1420–1428.
- Yilmaz AB (2003). Levels of heavy metals (Fe, Cu, Ni, Cr, Pb and Zn) in tissue of *Mugil cephalus* and *Trachurus mediterraneus* from Iskenderum Bay, Turkey. *Environmental Research*, 92(3): 277-281.